High Fiber Diet

- The ideal amount of daily fiber intake is 25-35 grams per day. Fiber draws in fluid from your body to add bulk to your stools and can make bowel movements soft or firm.
- Add fiber to your diet slowly. The bacteria in your stomach and small intestines need time to catch up. Adding too much fiber or adding fiber too quickly may cause gas, bloating, cramps and diarrhea. The MBCP recommends adding 5 grams of fiber to your diet every day at two week intervals. Initially there will be an increase in gas production however the gas production will decrease over time.
- Caffeine is a diuretic which pulls fluids from your body and excretes it as urine. Caffeine-free fluids allow more water to stay in your body. This allows the colon to absorb the fluid, making your bowel movements soft. For every 1 cup of caffeinated fluids, the MBCP recommends that you drink 2 glasses of decaffeinated fluids. Constipation may result if fiber rich diet is consumed without enough caffeine-free fluids.
- Most fiber supplements that are tablets only have 0.5 grams of fiber per tablet. Make sure to read the package for proper dosing. Few studies have been completed on the effectiveness of fiber supplements.
- Commonly used fibers to treat constipation include psyllium seed, synthetic cellulose (methyl cellulose), and calcium polycarbophil.
 Psyllium and methyl cellulose (and probably calcium polycarbophil) do not increase the production of gas; however, they still may result in the sensation of bloating.

Types of Fiber

	Insoluble Fiber	Soluble Fiber
Recommended		
Daily Amount	25-30 grams	3 grams
Functions	 Move bulk through the intestines Control and balance the pH (acidity) in the intestines 	 Absorb and retain water to form a bulking gel Bind with fatty acids Prolong stomach emptying time permitting a better absorption of nutrients
Benefits	 Promote regular bowel movements and prevent constipation. Bulk up stools that are too loose. Move toxic waste through colon in less time Help prevent colon cancer by keeping an optimal pH in intestines to prevent microbes from producing cancerous substances 	 Lower total cholesterol and LDL cholesterol (the bad cholesterol) therefore reducing the risk of heart disease Regulate blood sugar for people with diabetes

Examples of Caffeine-free Fluids:

- Water
- Milk
- Fruit and vegetable juices
- Caffeine-free pop
- Caffeine-free coffee/tea
- Flavored water
- Vitamin water, excluding Tropical Citrus Energy flavor

Amount of Fiber in Common Foods:

Beans	Serving	Fiber
Black-eyed Peas	½ cup	2.92
Brown Peas	½ cup	8.43
Green/String Beans	½ cup	2.23
Kidney Beans	½ cup	6.43
Lentils	½ cup	6.00
Lima Beans	½ cup	4.50
Navy Beans	½ cup	8.57
Northern Beans	½ cup	8.57
Pinto Beans	½ cup	8.48
Red Beans	½ cup	8.48
Wax/Yellow Beans	½ cup	1.62
White Beans	½ cup	8.57

Breads/Grains	Serving	Fiber
(check package)		
Bagel (most bagels)	1	1.5
Barley	½ cup	8.26
Black Bread	1 slice	1.72
Bran Bread	1 slice	2.09
Cornbread	1 med. Piece	1.60
Cornmeal	½ cup	5.17
Cracked Wheat	1 slice	1.66
English Muffin, Thomas	1	1.5
Flour:		
-Buckwheat	½ cup	3.14
-Rye	½ cup	5.22
-White	½ cup	2.01
-Whole Wheat	½ cup	5.34
French Bread	1 slice	0.5
Graham Crackers	1 square	0.24
Italian, Bakery Light	1 slice	2.5
Multigrain Bread	1 slice	1.51

Bowel Control Program High Fiber Diet - 3 -

Pancakes	1 slice	1.5
Pita, white	1 slice	1.0
Pumpernickel Bread	1 slice	1.72
Rye Bread	1 slice	0.40
Saltines	1 reg.	0.12
Seven Grain Bread, Bran'ola	1 slice	3.0
Wheat Bread, Bakery Light	1 slice	3.0
Wheat Germ	3 tbsp.	3.9
White Bread	1 slice	1.66
Whole Wheat Bread	1 slice	2.0

Cereals	Serving Fiber (gram	
All Bran	1 cup	1.0
Alpha-Bits	1/3 cup	4.0
Banana Nut Crunch	1/3 cup	13.0
Bran Buds, Kellogg's	1 cup	3.0
Cheerios	1 ounce	0.51
Corn Flakes, Kellogg's	¾ cup	0.0
Corn Pops	1 pack	5.6
Cracklin' Oat Bran,	1 packet	1.0
Kellogg's	½ cup	13.0
Cream of Wheat	1 cup	5.87
Fiber One, General Mills	1 ounce	1.5
Frosted Mini-Wheats	1 cup	2.0
Granola	1 pack	3.0
Honey Nut Cheerios,	1 cup	6.0
G'Mills	1 cup	2.0
Instant Oatmeal	½ cup	7.0
Kashi 7 Whole Grains	1 cup	8.0
Flakes	1 cup	10.0
Kashi 7 Whole Grains	¾ cup	5.0
Honey Puffs	1 cup	3.0
Kashi 7 Whole Grains	1 ounce	1.98
Nuggets	1 ounce	0.26

Bowel Control Program High Fiber Diet - 4 -

Kashi GoLean Crunch!	1 ounce	0.26	
Kashi GoLean Original	3 biscuits	1.14	
Kashi Heart to Heart	1 ounce	7.3	
Multi-Grain Cheerios	¾ cup	3.0	
Oat Bran, cooked	1 ounce	2.83	
Oat Flakes	1 ounce	4.05	
Post Shredded Wheat	2 biscuits	6.0	
Quaker Shredded Wheat	3 biscuits	7.0	
Raisin Bran, Kellogg's	1 cup	7.0	
Fruits	Serving	Fiber (grams)	
Apple with skip	1 medium	3.7	
Apple with skin	1 medium	3.7 2.4	
Apple without skin		2.4	
Applesauce	½ cup 3 medium		
Apricots		2.5	
Avocado	1 cup	7.3	
Banana	1 medium	2.7	
Blackberries	1 cup	7.6	
Blueberries (raw)	1 cup	4.0	
Boysenberries	1 cup	7.2	
Cantaloupe	1 cup pieces	1.3	
Cherries	10 cherries 1.3		
Fruit salad/cup	½ cup 1.3		
Grapefruit	½ medium 1.3		
Grapes	1 cup	1.2	
Honeydew Melon	1 cup pieces	1.0	
Mandarin oranges	½ cup	1.0	
Mango	1 cup	3.0	
Nectarine	1 medium	2.2	
Orange	1 medium	3.0	
Papaya	1 medium	5.5	
Peach	1 medium	1.7	
Pear	1 medium	4.0	
Pineapple	1 cup pieces	2.0	
Plum	1 medium	1.0	
Prunes (dried)	10 prunes	6.0	

Raisins (seedless)	2/3 cup	4.0	
Raspberries	1 cup	8.4	
Strawberries	1 cup	3.4	
Tangerine	1 medium	2.0	
Watermelon	1 cup pieces	0.8	
Vegetables	Serving	Fiber (grams)	
Artichoke, boiled	1 medium	6.2	
Asparagus, boiled	½ cup, 6 spears	1.4	
Baked Beans	1 cup	14.0	
Beets, boiled	1 cup	3.4	
Broccoli, boiled	½ cup	2.3	
Brussels Sprouts, boiled	½ cup	2.0	
Carrots	1 medium	2.0	
Cauliflower, boiled	½ cup	1.7	
Celery	1 stalk (7 inches)	0.7	
Coleslaw	½ cup	1.0	
Corn on the cob	1 ear	2.0	
Cucumber	½ cup of slices	0.5	
Eggplant, boiled	½ cup 1.0		
Green beans, boiled	½ cup	2.0	
Lima Beans, boiled	1 cup	13.2	
Lettuce	½ cup of pieces 0.5		
Mushrooms	½ cup of pieces 0.4		
Onions, boiled	½ cup 1.0		
Peas, Green	½ cup 4.0		
Pinto Beans, boiled	1 cup	14.7	
Potato, baked with skin	1 medium	5.0	
Potato, boiled	1 medium	2.0	
Potato salad	½ cup	1.6	
Pumpkin, canned	½ cup 5.0		
Rutabaga	1 cup	3.0	
Spinach, boiled	½ cup 2.2		
Spinach, raw	½ cup	0.8	
Squash, winter	½ cup	3.0	
Sweet Potato, baked	1 medium	3.0	
Tomato, raw	1 medium 1.0		

Bowel Control Program High Fiber Diet - 6 -

Turnip greens, cooked	1 cup 5.0		
Zucchini	½ cup	2.34	
Breads/Grains	Serving	Fiber (grams)	
(check package)			
Bagel (most bagels)	1	1.5	
English Muffin, Thomas	1	1.5	
French Bread	1 slice	0.5	
Italian, Bakery Light	1 slice	2.5	
Pancakes	1 slice	1.5	
Pita, white	1 16" diameter	1.0	
Seven grain, Bran'ola	1 slice	3.0	
Wheat, Bakery Light	1 slice	2.5	
Whole Wheat	1 slice	2.0	
Pasta	Serving	Fiber (grams)	
Elbow Macaroni, Golden	½ cup	2.0	
Grain	1 cup	1.8	
Macaroni	1 cup	4.0	
Macaroni, whole wheat	1 cup	6.3	
Spaghetti, whole wheat	1 cup	3.5	
Brown Rice, long	1 cup	1.0	
White rice	2 ounces 1.27		
Egg Noodles	2 ounces	7.0	
Ronzoni Smart Taste Pasta	2 ounces	7.0	
Nuts and Seeds	Serving	Fiber (grams)	
Almonds	10 nuts	1.1	
Brazil nuts	8 nuts	2.1	
Cashews	18 nuts	1.1	
Flaxseeds	1 tbsp.	3.3	
Peanuts	10 nuts	1.4	
Peanut butter	1 tbsp.	1.5	
Pecans	10 nuts	2.7	
Pistachios	50 nuts	2.9	
Sunflower seeds	1 tbsp.	0.5	
Walnuts	7 nuts	1.9	

Snacks	Serving	Fiber (grams)
Kashi TLC Cookies	1 cookie	4 g
Kashi GoLean Chewy Bars	1 bar	6 g
Kashi GoLean Crunchy!	1 bar	5 g
Bars		- 0
Kashi GoLean Roll! Bars	l bar	6 g
Kashi TLC Cereal Bars	l bar	3 g
Kashi TLC Chewy Granola	1 bar	4 g
Bars		C
Kashi TLC Crunchy	2 bars	4 g
Granola Bars		C
Quaker Breakfast Cookies	1 cookie	5 g
Quaker Chewy Bars	1 bar	1 g
Quaker Fiber & Omega-3	1 bar	9 g
bars		
Quaker Oatmeal to go High	1 bar	10 g
Fiber		
Fiber One Bar	1 bar	9 g
Fiber One Yogurt	1 container	5 g
Kellogg FiberPlus Bars	1 bar	9 g
LUNA Bars	1 bar	3 g

Foods Containing

Soluble Fiber:

- Rice
- Pasta and noodles
- Oatmeal
- Barley
- Fresh white breads such as french or sourdough
- Rice cereals
- Flour tortillas

- Soy
- Potatoes
- Carrots
- Yams
- Sweet potatoes
- Turnips
- Rutabagas
- Parsnips
- Beets

- Squash and pumpkins
- Mushrooms
- Chestnuts
- Avocados
- Bananas
- Applesauce
- Mangoes
- Papayas

Bowel Control Program High Fiber Diet - 8 -

Fiber Products

Brand	Type (serving size)	Amount of Fiber	Price
Acacia Fiber	Powder (1 tsp.)	2 g	\$16.99
Benefiber	Powder (2 tsp.) (types- regular or with B vitamins)	3 g (3 g soluble)	\$7.99
(wheat dextrin)	Caplets (3)	3 g (3 g soluble)	\$11.99
	Chewable (3)	3 g (3 g soluble)	\$11.99
	Caplets (3)	2 g (2 g soluble)	\$9.60
Citrucel	Chewable (3)	2 g (2 g soluble)	\$9.69
(methylcellulose)	Caplets (1)	0.5 g (0.5 g soluble)	\$14.32
Equalactin (calcium polycarbophil)	Tablets (2)	0.625 g	\$8.49
FiberChoice (inulin)	Chewable tablets (2)	4 g	\$14.99
FiberCon (calcium polycarbophil)	Caplet (1)	0.5 g (0.5 g soluble)	\$13.79
Hydrocil (psyllium)	Powder (1 tsp.)	3 g	\$13.99
Konsyl (psyllium)	Powder (1 tsp./6 g)	5 g (3 g soluble)	\$15.99
Metamucil	Powder (1 tsp./12 g) (Gluten-free)	3 g (2 g soluble)	\$8.59
(psyllium)	Wafers (2)	6 g (3 g soluble)	\$5.49
	Capsules (6) (Gluten-free)	3 g (0.33 g soluble)	\$12.51
Metamucil Clear & Natural (Fibersure) (inulin)	Powder (1 tsp.)	5 g (5 g soluble)	\$8.99
VitaFusion	Fiber Gummies (2 gummies)	5 g (5 g soluble)	\$13.99

Helpful Hints:

- Try a variety of brands or different forms of the same brand until you find the supplement that works for you.
- Avoid sugar-free supplements that contain artificial sweeteners as these can increase gas and bloating.

Psyllium

Psyllium found in fiber supplements is derived from the seeds of the Psyllium plant. The seeds expand and become gel-like when exposed to water in the body and add bulk to the stool to relieve either constipation or diarrhea. Psyllium is the most commonly found active ingredient in fiber products. Some individuals find it to cause excessive gas and bloating.

Methylcellulose

Methylcellulose found in fiber supplements is a synthetic product derived from cellulose. Methylcellulose is not broken down and digested in the intestines, but rather absorbs water and becomes gel-like to add bulk to the stool. Because methylcellulose does not ferment, it may cause less gas and bloating in some individuals.

Calcium Polycarbophil

Calcium Polycarbophil is a synthetic, bulk-forming fiber. It passes through the intestines undigested and absorbs water to add bulk to the stool, relieving either constipation or diarrhea.

Wheat Dextrin

Wheat Dextrin is a fiber derived from wheat starch. It expands and forms a gelatinous substance when exposed to water thus adding bulk to stool. Wheat dextrin may cause less gas and bloating than other fiber supplements in some individuals.

Inulin

Inulin is a compound produced by many plants and is composed mainly of fructose. Inulin is a prebiotic and, like other fiber supplements, it adds bulk to the stool to relieve constipation or diarrhea, but it may cause excessive gas and bloating in some individuals.

Acacia Fiber

Acacia Fiber is derived from the gum of the acacia tree. In addition to being a fiber supplement, it is also a prebiotic which promotes the growth of healthy bacteria in the gut for digestion. Acacia fiber adds bulk to the stool and does not cause excessive gas and bloating.

Gluten-free Fiber Supplements

- Citrucel
- Fibercon
- Konsyl
- Metamucil powder and capsules (not wafers)
- Acacia fiber

Disclaimer: This document is not intended to take the place of the care and attention of your personal physician or other professional medical services. Talk with your doctor if you have Questions about individual health concerns or specific treatment options.